



Comparison of metoprolol degradation by Fe^{III}-NTA modified Fenton-like reaction in the absence and presence of manganese: Efficiency and intermediates

Yifan Li^a, Jianhui Sun^a, Sheng-Peng Sun^{b,*}


^a Key Laboratory for Yellow River and Huai River Water Environmental and Pollution Control, Ministry of Education, Henan Key Laboratory for Environmental Pollution Control, School of Environment, Henan Normal University, Xinxiang, Henan 453007, PR China

^b Suzhou Key Laboratory of Green Chemical Engineering, School of Chemical and Environmental Engineering, College of Chemistry, Chemical Engineering and Materials Science, Soochow University, Suzhou, Jiangsu 215123, PR China

HIGHLIGHTS

- Metoprolol degradation by Fe^{III}-NTA/H₂O₂ was enhanced in the presence of Mn^{II}.
- Rapid degradation of metoprolol was obtained by Fe^{III}-NTA/H₂O₂-Mn at pH 4.0–8.0.
- Fe^{III}-NTA/H₂O₂-Mn led to more efficient degradation of metoprolol and intermediates.
- Degradation pathways of metoprolol were proposed.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 7 October 2016
Received in revised form 21 December 2016
Accepted 22 December 2016
Available online 26 December 2016

Keywords:

Metoprolol
Fe^{III}-NTA
Fenton-like reaction
Hydroxyl radical
Manganese

ABSTRACT

The degradation of metoprolol in aqueous solutions by traditional Fenton-like reaction (Fe^{III}/H₂O₂), and Fe^{III}-NTA modified Fenton-like reaction (Fe^{III}-NTA/H₂O₂) in the absence and presence of Mn^{II} (Fe^{III}-NTA/H₂O₂-Mn) have been investigated. The results show that Fe^{III}-NTA/H₂O₂ is able to degrade metoprolol at initial neutral pH. In particular, the presence of Mn^{II} greatly improved the degradation rate of metoprolol by Fe^{III}-NTA/H₂O₂ over a wide pH range of 4.0–8.0. Under the same conditions, the degradation rate constants of metoprolol (*k*) obtained in Fe^{III}-NTA/H₂O₂-Mn system were typically 7–9-fold larger than those obtained in Fe^{III}-NTA/H₂O₂ system. The involved reactions in Fe^{III}-NTA/H₂O₂-Mn system were proposed on the basis of important parameters analysis including ferrous ion concentration, dissolved oxygen (DO) concentration, and the quenching experiments for hydroxyl radical (HO·) and superoxide anion radical (O₂^{•−}). Several intermediates have been identified by mass spectrometry. Our results suggest that the degradation of metoprolol in both Fe^{III}-NTA/H₂O₂ and Fe^{III}-NTA/H₂O₂-Mn systems were caused by HO· attack. The degradation pathways of metoprolol were proposed on the basis of the identified intermediates. Fe^{III}-NTA/H₂O₂-Mn system led to more efficient degradation of metoprolol and its intermediates.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

Metoprolol, a selective β₁ receptor blocking drug, is widely used in the therapy of hypertension and angina. In the past decade, metoprolol and its metabolites have been detected in sewage treatment plant effluents, surface water, groundwater, and even

* Corresponding author.

E-mail address: shepsun@suda.edu.cn (S.-P. Sun).

in drinking water, which pose a potential risk to human health [1,2]. The degradation rate of metoprolol in water by the direct photolysis is small, e.g., the extrapolated half-lives of metoprolol in central Europe and the U.S. were reported at 28–95 d in summer and 190–449 d in winter [3]. Thus, metoprolol is most likely to be pseudo-persistent in natural waters if do not consider the indirect photolysis effects. In addition, the efficiency of metoprolol removal is lower than 30% in traditional sewage treatment plant [4]. There is an urgent need to improve the removal efficiencies of pharmaceuticals in water.

A far more effective method is the so-called advanced oxidation processes (AOPs), involving the generation of hydroxyl radical ($\text{HO}^\bullet$) [5–7], have been widely applied in water treatment for harmful organic compounds removal. Efficient degradation of metoprolol has been obtained by photocatalysis, UV/ H_2O_2 and Fenton-related processes [8–10]. Two decades ago, research by Sun and Pignatello [11] showed the successful degradation of 2,4-dichlorophen-oxyacetic acid at pH 6.0 by chelates modified Fenton-like reaction. Chelating agents modified Fenton-like reaction has been receiving increasing attention because it can work at neutral pH conditions, which overcomes the limitation of acidic pH (e.g., optimal pH 2.8–3.0) by using traditional Fenton and/or Fenton-like reaction. S,S-ethylenediamine-N,N'-disuccinic acid (EDDS) modified Fenton reaction (i.e., Fe^{II} -EDDS/ H_2O_2) and Fenton-like reaction (i.e., Fe^{III} -EDDS/ H_2O_2) are capable of degrading 4-chlorophenol, bisphenol A, naphthenic acid, and 4-tert-butylphenol at near neutral or basic pH conditions [12–15]. The use of Fe^{III} -EDDS for photo-Fenton reaction at neutral pH is also successful in degrading micro-pollutants in wastewater treatment plant effluents [16–18]. In addition, nitrilotriacetic acid (NTA) and ethylenediaminetetraacetic acid (EDTA) modified Fenton-like reactions (i.e., Fe^{III} -NTA/ H_2O_2 and Fe^{III} -EDTA/ H_2O_2) have also been proved to be highly effective in degrading atrazine, carbamazepine, fenuron, naphthenic acid, parachlorobenzoic acid and sulfamethoxazole at neutral or slightly basic pH conditions [19–22]. In particular, our recent finding suggested that the presence of manganese (Mn^{II}) could greatly enhance the Fenton-like catalytic activity of Fe^{III} -NTA/ H_2O_2 [23]. The potential mechanism may involve an enhanced generation of superoxide anion radical ($\text{O}_2^{\bullet-}$) by the Mn^{II} , which can accelerate the reduction of Fe^{III} -NTA to Fe^{II} -NTA and indirectly improve the generation of $\text{HO}^\bullet$ [23]. The Mn^{II} / Fe^{III} -NTA catalytic system would be very promising since Fe^{III} and Mn^{II} are quite ubiquitous in the environment and can be used as natural Fenton-like catalyst in the presence of NTA (or natural Fe^{III} complexing agents) and H_2O_2 . However, little research on this system has been reported.

The purpose of this study is to demonstrate the potential effectiveness of Fe^{III} -NTA/ H_2O_2 -Mn for water treatment. In the present study, we choose metoprolol as a target compound, to evaluate the efficiency and intermediates generated from metoprolol degradation by Fe^{III} -NTA/ H_2O_2 in the absence and presence of Mn^{II} . The degradation kinetics of metoprolol in both systems were compared under various conditions. The involved reactions in Fe^{III} -NTA/ H_2O_2 -Mn system were proposed on the basis of certain important parameters analysis such as ferrous ion concentration, dissolved oxygen (DO) concentration, and the quenching experiments for $\text{HO}^\bullet$ and $\text{O}_2^{\bullet-}$. The intermediates were identified by liquid chromatography/mass spectrometry.

2. Materials and methods

2.1. Chemicals

Ferric chloride hexahydrate, hydrogen peroxide (30% H_2O_2 , w/w), sodium hydroxide and sulfuric acid were purchased from Sino-

pharm Chemical Reagent Co., Ltd. Metoprolol tartrate (purity $\geq 98\%$) was purchased from LKT Laboratories, Inc. NTA was purchased from Acros Organics. Formic acid and manganese nitrate tetrahydrate were purchased from Sigma-Aldrich. HPLC-grade methanol, 2-propanol and water were purchased from Fisher Scientific. Chloroform was purchased from Chinasun Specialty Products Co., Ltd. All chemicals were of reagent grade or better, and used as received without further purification. Milli-Q water was used to prepare solutions.

2.2. Experimental procedures

All experiments were performed in 125 ml glass Erlenmeyer flasks. Briefly, 100 ml of metoprolol aqueous solution and an appropriate amount of freshly prepared Fe^{III} -NTA solution were added into the flask. The flask was then placed in a water-jacketed glass vessel water bath ($25 \pm 1^\circ\text{C}$) under a magnetic stirring condition. After that, an appropriate amount of Mn^{II} was added to the solution. The pH of the solution was then adjusted to the desired value using 1.0/0.1 M NaOH or 1.0/0.1 M H_2SO_4 . Finally, Fenton-like reaction was started by the addition of an appropriate amount of H_2O_2 . It is worth noting that the water-jacketed glass vessel was covered with aluminum foil to prevent the effect of light on the Fenton-like reaction. Samples (1 ml) were withdrawn at different time intervals and mixed with 0.1 ml of methanol immediately to quench the reaction.

2.3. Analytical methods

The concentration of metoprolol was determined by a high performance liquid chromatography (HPLC, Agilent 1100) with fluorescence detection. An Ultra C-18 column ($5\ \mu\text{m}$, $250 \times 4.6\ \text{mm}$) was used for the chromatographic separation. The mobile phase consisted of methanol and 0.1% formic acid aqueous solution (35:65, v/v) at a flow rate of $0.9\ \text{ml min}^{-1}$. The column temperature was 30°C . The injection volume was $20\ \mu\text{l}$, and the fluorescence detection was performed at 275 nm of excitation and 298 nm of emission.

The degradation intermediates were identified by an Agilent 1200 HPLC system coupled to a 6130 quadrupole mass spectrometer. The MS detector was performed for a full scan range of m/z 30–600 under an ESI + APCI positive mode. The MS operating conditions were as follows: Fragmentor 90, drying gas flow rate $6\ \text{L min}^{-1}$, nebulizer pressure 45 psi, dry gas temperature 300°C , vaporizer temperature 250°C , capillary voltage 3500 V, corona current $4.0\ \mu\text{A}$ and charging voltage 2000 V.

The concentration of ferrous ion in the aqueous solutions was determined by using the 1,10-phenanthroline method. The concentration of H_2O_2 was determined by the iodide method [24]. The dissolved oxygen (DO) concentration in the aqueous solutions during the reaction time was monitored by a dO_2 sensor (Model Z010023525, AppliSens).

3. Results and discussion

3.1. Comparison of metoprolol degradation by Fe^{III} / H_2O_2 , Fe^{III} -NTA/ H_2O_2 and Fe^{III} -NTA/ H_2O_2 -Mn

Fig. 1 shows a comparison of metoprolol degradation by Fe^{III} / H_2O_2 , Fe^{III} -NTA/ H_2O_2 and Fe^{III} -NTA/ H_2O_2 -Mn systems at initial neutral pH. A minimum degradation of metoprolol was observed in Fe^{III} / H_2O_2 system, with 3.61% degradation efficiency of metoprolol after 60 min of the reaction time. This result was expected because of the precipitation of ferric ion at pH 7.0 in the absence of chelator. In contrast, Fe^{III} -NTA complex is able to catalyze




Fig. 1. Comparison of metoprolol degradation by $\text{Fe}^{\text{III}}/\text{H}_2\text{O}_2$, $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ and $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ systems. Experimental conditions: $29.2\ \mu\text{M}$ of metoprolol, $0.1\ \text{mM}$ of Fe^{III} , $\text{NTA}/\text{Fe}^{\text{III}}$ molar ratio of 2:1, $\text{H}_2\text{O}_2/\text{Fe}^{\text{III}}$ molar ratio of 100:1, $\text{Mn}^{\text{II}}/\text{Fe}^{\text{III}}$ molar ratio of 1:1, and initial pH of 7.0.

Fenton-like reaction at pH 7.0, resulted in 100% degradation efficiency of metoprolol after 30 min of the reaction time in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ system. In particular, $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system led to a very fast degradation of metoprolol, with 100% degradation efficiency of metoprolol after 7 min of the reaction time. The result is consistent with our recent finding that the presence of Mn^{II} could enhance $\text{Fe}^{\text{III}}\text{-NTA}$ -catalyzed Fenton-like reaction [23].

The effects of Mn^{II} concentration on the degradation kinetics of metoprolol in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system were shown in Fig. 2. An increase in Mn^{II} concentration from 0.01 to 0.1 mM, corresponding to $\text{Mn}:\text{Fe}$ molar ratios of 0.1:1 to 1:1, had a positive effect on metoprolol degradation. The apparent degradation rate constant of metoprolol (k) increased from 0.7547 to $1.6048\ \text{min}^{-1}$. However, further increasing Mn^{II} concentration from 0.2 to 1.0 mM led to a slight inhibitory effect on the degradation rate of metoprolol, e.g., the k value of metoprolol decreased from 1.6289 to $1.1103\ \text{min}^{-1}$. This is because when Mn^{II} concentration was larger than 0.2 mM, hydrous manganese dioxide could form from free Mn^{II} in excess under the given conditions, which is capable of catalyzing the decomposition of H_2O_2 to O_2 and H_2O [25]. It is worth noting that even at a high concentration of 1.0 mM of Mn^{II} , corresponding to $\text{Mn}:\text{Fe}$ molar ratio of 10:1, the k value of metoprolol was still larger (6-fold) than that in the absence of Mn^{II} .

Quenching tests showed that the degradation of metoprolol in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system was significantly inhibited by the addi-




Fig. 2. Effects of Mn^{II} concentration on the degradation kinetics of metoprolol in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system. Experimental conditions: $29.2\ \mu\text{M}$ of metoprolol, $0.1\ \text{mM}$ of Fe^{III} , $\text{NTA}/\text{Fe}^{\text{III}}$ molar ratio of 2:1, $\text{H}_2\text{O}_2/\text{Fe}^{\text{III}}$ molar ratio of 100:1, $\text{Mn}^{\text{II}}/\text{Fe}^{\text{III}}$ molar ratio of 0.1:1–10:1, and initial pH of 7.0.

tion of 2-propanol, but not by chloroform (Fig. S1, in Supplementary data), which suggests that the degradation of metoprolol was probably caused by $\text{HO}^\bullet$ attack. The presence of Mn^{II} is unable to directly catalyze Fenton-like reaction to generate $\text{HO}^\bullet$; this is because no degradation of metoprolol was observed in the absence of Fe^{III} (e.g., $\text{Mn}^{\text{II}}\text{-NTA}/\text{H}_2\text{O}_2$ in Fig. 1). Thus, the very fast degradation of metoprolol in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system could be attributed to an indirect enhancement of $\text{Fe}^{\text{III}}\text{-NTA}$ -catalyzed Fenton-like reaction by Mn^{II} . One hypothesis is that the presence of Mn^{II} favors the production of $\text{O}_2^{\cdot-}$, which enhances the reduction of $\text{Fe}^{\text{III}}\text{-NTA}$ to $\text{Fe}^{\text{II}}\text{-NTA}$, and in turn increases the production of $\text{HO}^\bullet$ from the reaction of $\text{Fe}^{\text{II}}\text{-NTA}$ with H_2O_2 . In order to prove this, the concentration of ferrous ion in the aqueous solutions was monitored during the reaction time, and the results were shown in Fig. 3. It can be seen that the concentration of ferrous ion in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system increased rapidly from 0 to $0.40\ \text{mg L}^{-1}$ within 5 min of reaction time, and then gradually decreased to $0.11\ \text{mg L}^{-1}$ within 30 min of reaction time. On the contrary, the concentration of ferrous ion in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ system was always below $0.02\ \text{mg L}^{-1}$, which was much smaller than that in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system. In addition, changes in DO concentrations in both systems with the reaction time were also monitored. It can be observed from Fig. 3 that the DO concentration in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system increased significantly from 100% (i.e., 100% air saturation) to 275% within 7 min of reaction time, and then gradually decreased to 211% within 30 min of reaction time, which was much higher than that in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ system. The results suggested that more oxygen was produced in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system. Moreover, Fig. 3 provides further evidence that the decomposition rate of H_2O_2 in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system was faster than that in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ system.

On the basis of Haber-Weiss reactions, the involved main reactions in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system were proposed by R1–R8. As shown in reactions R1–R3, it was $\text{O}_2^{\cdot-}$ and not free $\text{HO}^\bullet$ which was generated from the reaction of $\text{Mn}^{\text{II}}\text{-NTA}$ with H_2O_2 . On the basis of reactions R4 and R5, $\text{O}_2^{\cdot-}$ should rapidly reduce $\text{Fe}^{\text{III}}\text{-NTA}$ (and $\text{Mn}^{\text{III}}\text{-NTA}$) to $\text{Fe}^{\text{II}}\text{-NTA}$ (and $\text{Mn}^{\text{II}}\text{-NTA}$), leading to the production of oxygen. Thus, R1–R5 also explain the rapid decomposition of H_2O_2 and the significant increase of DO concentration in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system. Most importantly, $\text{HO}^\bullet$ is generated from the reaction of $\text{Fe}^{\text{II}}\text{-NTA}$ with H_2O_2 , which lead to metoprolol destruction (as shown in R6 and R8). Because $\text{Fe}^{\text{II}}\text{-NTA}$ is mainly generated










Fig. 3. Changes in DO concentration, H_2O_2 concentration and ferrous ion concentration with the reaction time in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ and $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ systems. Experimental conditions: $29.2\ \mu\text{M}$ of metoprolol, $0.1\ \text{mM}$ of Fe^{III} , $\text{NTA}/\text{Fe}^{\text{III}}$ molar ratio of 2:1, $\text{H}_2\text{O}_2/\text{Fe}^{\text{III}}$ molar ratio of 100:1, $\text{Mn}^{\text{II}}/\text{Fe}^{\text{III}}$ molar ratio of 1:1, and initial pH of 7.0.

from the reaction of $\text{Fe}^{\text{III}}\text{-NTA}$ with H_2O_2 in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ system (i.e., R7), thus the generation rate of $\text{Fe}^{\text{II}}\text{-NTA}$ in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ system is much slower than that in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system.



3.2. Effects of initial pH

The effects of initial pH on the degradation kinetics of metoprolol by $\text{Fe}^{\text{III}}/\text{H}_2\text{O}_2$, $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ and $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ systems were investigated over the range of pH 3.0–9.0 (Fig. 4). In $\text{Fe}^{\text{III}}/\text{H}_2\text{O}_2$ system, a k value of 0.0976 min^{-1} was obtained at an initial pH 3.0. When the initial pH value was higher than 5.0, the degradation of metoprolol was significantly inhibited in $\text{Fe}^{\text{III}}/\text{H}_2\text{O}_2$ system due to the precipitation of ferric ion. In $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ system, an increase in the initial pH value from 3.0 to 9.0 had a positive effect on the degradation rate of metoprolol, resulted in increasing the k value of metoprolol from 0.0033 to 0.7190 min^{-1} . The small degradation rates of metoprolol under acidic pH conditions (e.g., pH 3.0–4.0) can be explained either by relatively low regeneration rate of $\text{Fe}^{\text{II}}\text{-NTA}$, or by low catalytic activity of $\text{Fe}^{\text{III}}\text{-NTA}$ species toward H_2O_2 at acidic pH. First of all, $\text{O}_2^{\cdot-}$ was believed to play an important role in reducing $\text{Fe}^{\text{III}}\text{-NTA}$ to $\text{Fe}^{\text{II}}\text{-NTA}$ which then react with H_2O_2 to generate $\text{HO}\cdot$. Since more $\text{O}_2^{\cdot-}$ changes to perhydroxyl radical ($\text{HOO}\cdot$) at pH 3.0–4.0 (Eq. (1)) [26], the regeneration rate of $\text{Fe}^{\text{II}}\text{-NTA}$ can be limited due to the weak reduction capability of $\text{HOO}\cdot$.



On the other hand, $\text{Fe}^{\text{III}}\text{-NTA}$ species and free NTA also react with $\text{HO}\cdot$. The reaction between $\text{HO}\cdot$ and $\text{Fe}^{\text{III}}\text{-NTA}$ species/free




Fig. 4. Effects of initial pH on the degradation kinetics of metoprolol by $\text{Fe}^{\text{III}}/\text{H}_2\text{O}_2$, $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ and $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ systems. Experimental conditions: 29.2 μM of metoprolol, 0.1 mM of Fe^{III} , NTA/ Fe^{III} molar ratio of 2:1, $\text{H}_2\text{O}_2/\text{Fe}^{\text{III}}$ molar ratio of 100:1, $\text{Mn}^{\text{II}}/\text{Fe}^{\text{III}}$ molar ratio of 1:1, and initial pH of 3.0–9.0.

NTA would have an inhibiting effect on the degradation rate of metoprolol since the reactions are competing. A larger reaction rate between $\text{HO}\cdot$ and $\text{Fe}^{\text{III}}\text{-NTA}$ species/free NTA would lead to a smaller degradation rate of metoprolol. According to the literature [21,27], $\text{Fe}^{\text{III}}\text{-NTA}$ species and free NTA are strongly pH-dependent. Although the reaction rate constant of $\text{Fe}^{\text{III}}\text{-NTA}$ species with $\text{HO}\cdot$ is rarely reported (e.g., $1.6 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ was reported at pH 2.0 [28]), the reaction rate constants of $\text{HO}\cdot$ with free NTA have been reported at 6.1×10^7 , 7.5×10^8 , 5.5×10^8 , 4.77×10^8 , 2.5×10^9 and $4.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ at pH 2.0, 4.0, 6.0, 8.0, 9.0 and 10 [21,27], respectively. It can be seen that the reaction rate of $\text{HO}\cdot$ with free NTA was increased with the increase of pH from 2.0 to 10. Thus, the decrease of pH should favor the degradation of metoprolol. However, the present and previous studies have shown that the fast degradation of model contaminants and the decomposition of H_2O_2 occurred at neutral or slight basic pH in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ system [20,27], which rules out the effect of free NTA. Therefore, it is logically possible that $\text{Fe}^{\text{III}}\text{-NTA}$ species at acidic pH (i.e., $\text{Fe}^{\text{III}}(\text{NTA})(\text{H}_2\text{O})_2$ [29]) might be inefficient for the activation of H_2O_2 to induce Fenton-like reaction, or the reaction of $\text{Fe}^{\text{III}}(\text{NTA})(\text{H}_2\text{O})_2$ with $\text{HO}\cdot$ might act as an inhibitory reaction.

Maximum degradation rates of metoprolol were obtained in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system. It can be seen that the k values of metoprolol were much larger than those in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ system over the initial pH range of 4.0–8.0. At initial pH 3.0, the k value of metoprolol was 0.0819 min^{-1} in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system, which was close to that in $\text{Fe}^{\text{III}}/\text{H}_2\text{O}_2$ system (0.0976 min^{-1}). With increasing the initial pH from 3.0 to 5.0, the k values of metoprolol increased greatly from 0.0819 to 2.5809 min^{-1} in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system. On the basis of the pK_a value of $\text{HOO}\cdot/\text{O}_2^{\cdot-}$ ($\text{pK}_a = 4.8$) [26], the fraction of $\text{O}_2^{\cdot-}$ was calculated at 1.56%, 13.68% and 61.31% (of the total amount of $\text{HOO}\cdot$ and $\text{O}_2^{\cdot-}$) at pH 3.0, 4.0 and 5.0, respectively. It can be seen that the increase of the k values of metoprolol was consistent with the increase of the fraction of $\text{O}_2^{\cdot-}$ at pH 3.0–5.0. Thus, the results support our hypothesis that $\text{O}_2^{\cdot-}$ plays an important role in reducing $\text{Fe}^{\text{III}}\text{-NTA}$ to $\text{Fe}^{\text{II}}\text{-NTA}$, which in turn improves the generation of $\text{HO}\cdot$ and leads to a rapid degradation of metoprolol. It is worth noting that the fraction of $\text{HOO}\cdot$ is 38.69% at pH 5.0. $\text{HOO}\cdot$ is also an oxidant with an oxidation potential of 1.78 V [30], weaker than $\text{HO}\cdot$, thus the fastest degradation of metoprolol at pH 5.0 can be explained by $\text{HOO}\cdot$ along with $\text{HO}\cdot$ attack. At neutral pH, the fraction of $\text{O}_2^{\cdot-}$ would increase to >99% and the fraction of $\text{HOO}\cdot$ would decrease to <1%. Therefore, the degradation of metoprolol at neutral pH would be mainly caused by $\text{HO}\cdot$ attack. The large fraction of $\text{O}_2^{\cdot-}$ at neutral pH or slight basic pH could favor the generation of $\text{Fe}^{\text{II}}\text{-NTA}$. Unlike $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ system, however, the k values of metoprolol in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system gradually decreased with the increasing initial pH from 5.0 to 9.0. In particular, at initial pH 9.0, the k value of metoprolol in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ system was smaller than that in $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ system. This can be explained by the fact that the complexation constants of NTA with $\text{Fe}^{\text{III}}/\text{Fe}^{\text{II}}$ ($-\log k = 15.9$ and 9.8) are much larger than that with Mn^{II} ($-\log k = 7.46$) [31], thus Mn^{II} would firstly leach due to the degradation of NTA by $\text{HO}\cdot$ attack. The leached Mn^{II} could form hydrous manganese dioxide when the solution pH value is >5.0, which can directly catalyze the decomposition of H_2O_2 to O_2 and H_2O [25]. Additionally, the decrease of Mn^{II} concentration in aqueous phase would lead to the rate of $\text{O}_2^{\cdot-}$ production decreasing, which in turn decrease the generation rates of $\text{Fe}^{\text{II}}\text{-NTA}$ and $\text{HO}\cdot$.

3.3. Effects of NTA:Fe molar ratios

The effects of NTA/ Fe molar ratios on the degradation kinetics of metoprolol by $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2$ and $\text{Fe}^{\text{III}}\text{-NTA}/\text{H}_2\text{O}_2\text{-Mn}$ systems at initial neutral pH were investigated (Fig. 5). It can be seen that over

the range of NTA:Fe molar ratios tested, the k values of metoprolol in Fe^{III} -NTA/ H_2O_2 -Mn system were 7–17-fold larger than those in Fe^{III} -NTA/ H_2O_2 system. An increase in NTA/Fe molar ratio from 0.5:1 to 2:1 had positive effects on the degradation rates of metoprolol in both Fe^{III} -NTA/ H_2O_2 and Fe^{III} -NTA/ H_2O_2 -Mn systems. The k values of metoprolol increased from 0.0311 to 0.1754 min^{-1} in Fe^{III} -NTA/ H_2O_2 system and increased from 0.3450 to 1.6048 min^{-1} in Fe^{III} -NTA/ H_2O_2 -Mn system, respectively. When NTA/Fe molar ratio was larger than 2:1, however, a slight decrease in the k values of metoprolol was observed in either Fe^{III} -NTA/ H_2O_2 or Fe^{III} -NTA/ H_2O_2 -Mn system. This can be explained by the fact that the free NTA in excess scavenges $\text{HO}^\cdot$. Although the reaction rate constant of metoprolol with $\text{HO}^\cdot$ ($5.2\text{--}8.4 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$) [32,33] is larger than that of Fe^{III} -NTA/free NTA with $\text{HO}^\cdot$, it is believed that the competitive consumption of $\text{HO}^\cdot$ by the free NTA in excess can slightly limit the degradation of metoprolol.

3.4. Effects of H_2O_2 : Fe^{III} molar ratios

Fig. 6 shows the effects of H_2O_2 / Fe^{III} molar ratios on the degradation kinetics of metoprolol by Fe^{III} -NTA/ H_2O_2 and Fe^{III} -NTA/ H_2O_2 -Mn systems at initial neutral pH. Without the addition of H_2O_2 , no degradation of metoprolol was observed in both systems. Over the range of H_2O_2 / Fe^{III} molar ratios tested, the k values of metoprolol in Fe^{III} -NTA/ H_2O_2 -Mn system were 5–9-fold larger than those in Fe^{III} -NTA/ H_2O_2 system. With increasing H_2O_2 / Fe^{III} molar ratio from 10:1 to 200:1, the k values of metoprolol continuously increased from 0.0598 to 0.2276 min^{-1} in Fe^{III} -NTA/ H_2O_2 system. In Fe^{III} -NTA/ H_2O_2 -Mn system, the k values of metoprolol increased greatly from 0.3384 to 1.6048 min^{-1} and reached a plateau with increasing H_2O_2 / Fe^{III} molar ratio from 100:1 to 200:1. It is expected that the use of higher molar ratio of H_2O_2 / Fe^{III} led to higher rate of $\text{HO}^\cdot$ generation, by which faster degradation of metoprolol could be obtained. No inhibitory effect of H_2O_2 on the degradation kinetics of metoprolol was observed even at H_2O_2 / Fe^{III} molar ratio of 200:1. This is because the reaction rate constant of H_2O_2 with $\text{HO}^\cdot$ ($(1.2\text{--}4.5) \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$) [34] is much smaller than those of metoprolol, Fe^{III} -NTA and free NTA with $\text{HO}^\cdot$.

3.5. Effect of Fe^{III} dosage

Fe^{III} acts as the catalyst to initiate Fenton-like reaction, a high rate of $\text{HO}^\cdot$ generation might be obtained at high concentration of Fe^{III} -NTA. The effects of Fe^{III} dosages on the degradation kinetics of metoprolol by Fe^{III} -NTA/ H_2O_2 and Fe^{III} -NTA/ H_2O_2 -Mn systems at




Fig. 5. Effects of NTA/ Fe^{III} molar ratios on the degradation kinetics of metoprolol by Fe^{III} -NTA/ H_2O_2 and Fe^{III} -NTA/ H_2O_2 -Mn systems. Experimental conditions: 29.2 μM of metoprolol, 0.1 mM of Fe^{III} , NTA/ Fe^{III} molar ratio of 0.5:1–5:1, H_2O_2 / Fe^{III} molar ratio of 100:1, Mn^{II} / Fe^{III} molar ratio of 1:1, and initial pH of 7.0.




Fig. 6. Effects of H_2O_2 / Fe^{III} molar ratios on the degradation kinetics of metoprolol by Fe^{III} -NTA/ H_2O_2 and Fe^{III} -NTA/ H_2O_2 -Mn systems. Experimental conditions: 29.2 μM of metoprolol, 0.1 mM of Fe^{III} , NTA/ Fe^{III} molar ratio of 2:1, H_2O_2 / Fe^{III} molar ratio of 5:1–200:1, Mn^{II} / Fe^{III} molar ratio of 1:1, and initial pH of 7.0.




Fig. 7. Effects of Fe^{III} dosages on the degradation kinetics of metoprolol by Fe^{III} -NTA/ H_2O_2 and Fe^{III} -NTA/ H_2O_2 -Mn systems. Experimental conditions: 29.2 μM of metoprolol, 0.01–0.15 mM of Fe^{III} , NTA/ Fe^{III} molar ratio of 2:1, H_2O_2 / Fe^{III} molar ratio of 100:1, Mn^{II} / Fe^{III} molar ratio of 1:1, and initial pH of 7.0.





Fig. 8. Changes in the total molecular ion chromatogram area of metoprolol and its intermediates with the reaction time. Experimental conditions: 29.2 μM of metoprolol, 0.1 mM of Fe^{III} , NTA/ Fe^{III} molar ratio of 2:1, H_2O_2 / Fe^{III} molar ratio of 100:1, Mn^{II} / Fe^{III} molar ratio of 1:1, and initial pH of 7.0.

initial neutral pH were shown in Fig. 7. No degradation of metoprolol was observed in the absence of Fe^{III} , which suggests that $\text{HO}^\cdot$ is generated from the Fenton-like reaction of Fe^{III} -NTA/ H_2O_2 . Over



Scheme 1. Intermediates and degradation pathways of metoprolol.

the range of Fe^{III} concentrations tested, the k values of metoprolol in Fe^{III} -NTA/ H_2O_2 -Mn system were 7–9-fold larger than that in Fe^{III} -NTA/ H_2O_2 system. An increase in Fe^{III} concentrations from 0 to 0.15 mM has positive effects on the degradation rate of metoprolol in both systems. The k values of metoprolol increased from

0 to 0.3174 min^{-1} in Fe^{III} -NTA/ H_2O_2 system and to 2.2988 min^{-1} in Fe^{III} -NTA/ H_2O_2 -Mn system, respectively. The results indicate that the rate of $\text{HO}^\bullet$ generation is greatly enhanced by increasing Fe^{III} -NTA concentration, especially in Fe^{III} -NTA/ H_2O_2 -Mn system. This can be explained by the fact that high concentration of

Fe^{III}-NTA would increase the generation rate of Fe^{II}-NTA (R5) and (R7), which in turn reacts with H₂O₂ to produce more HO[•].

3.6. Intermediates and degradation pathways

It was worth noting that a high initial concentration of metoprolol (292 μM, ten times higher than that in typical experiments) was used in order to capture more intermediates. The typical full-scan total molecular ion chromatograms of samples at different reaction time (RT) can be found in the [Supplementary data](#). Two major peaks with molecular ion ([M+H]⁺) at mass-to-charge (*m/z*) ratios of 268 and 192 were observed in samples before the treatment, corresponding to metoprolol and NTA, respectively. Several intermediates were detected in samples after the reaction by mass spectrometry, their molecular ion were identified at *m/z* 134, 238, 258, 274, 282, 284, 298, 300, 314, 316, 330, 332 and 348. A major degradation product is 3-(isopropylamino)propane-1,2-diol (*m/z* 134), which formed by the cleavage of aromatic ether bond of metoprolol due to the attack of HO[•]. A same intermediate has been detected in the degradation of metoprolol by photocatalysis, γ-irradiation, Fenton and photo-Fenton processes [8,10,33,35]. In addition, the attack of HO[•] on the ether bond and/or the β-C in methoxyethyl side chain of metoprolol gave intermediates with molecular ion at *m/z* 238, corresponding to 4-(2-hydroxy-3-(isopropylamino)propoxy)benzaldehyde and 1-(4-ethylphenoxy)-3-(isopropylamino)propan-2-ol, consistent with previous studies [8,10].

The hydroxylation of benzene ring (and/or side chain) of metoprolol led to the formation of hydroxyl-metoprolol isomers (*m/z* 284). The intermediates with molecular ion at *m/z* 282 could be 4-(2-methoxyethyl)phenyl 2-hydroxy-3-(isopropylamino)propanoate and 1-(4-(2-hydroxy-3-(isopropylamino)propoxy)phenyl)-2-methoxyethanone, which were formed by the oxidation of hydroxyl group to keto group in hydroxyl-metoprolol (i.e., 3-(isopropylamino)-1-(4-(2-methoxyethyl)phenoxy)propane-1,2-diol and 1-(4-(1-hydroxy-2-methoxyethyl)phenoxy)-3-(isopropylamino)propan-2-ol). Moreover, the addition of 1 HO[•], 2 HO[•], 3 HO[•] and 4 HO[•] on hydroxyl-metoprolol gave intermediates with molecular ion at *m/z* 300, 316, 332 and 348, corresponding to di-, tri-, tetra-, and penta-hydroxyl-metoprolol, respectively. The intermediates with molecular ion at *m/z* 298 and 314 could be formed by the addition of 1 HO[•] and 2 HO[•] on the detected intermediates (*m/z* 282). The intermediates with molecular ion at *m/z* 258 and 274 were probably formed by the addition of 2 HO[•] and 3 HO[•] on two undetected intermediates (*m/z* 226), i.e., 1-amino-3-(4-(2-methoxyethyl)phenoxy)propan-2-ol and 4-(2-hydroxy-3-(isopropylamino)propoxy)phenol [33,36].

A comparison of Figs. S2 and S3 demonstrates that Fe^{III}-NTA/H₂O₂-Mn system led to more efficient degradation of metoprolol and its intermediates. In addition, Fig. 8 shows the changes in the total molecular ion chromatogram area of metoprolol and its intermediates with the reaction time. It can be seen that the total areas of major intermediates (i.e., *m/z* 238, 282, 284, 298, 300, etc.) increased rapidly to reach their maximum values within the first 3–5 min of the reaction time, and then decreased to undetectable levels at 20 min of the reaction time. In particular, the major intermediate (*m/z* 134) increased continuously until almost complete degradation of metoprolol took place (12–15 min of the reaction time), and then began to decrease slowly over time. The degradation pathways of metoprolol were proposed on the basis of these identified intermediates, which were presented in Scheme 1. Our results clearly suggested that the degradation of metoprolol and its intermediates were mainly caused by HO[•] attack.

4. Conclusion

The results of this study indicate that Fe^{III}-NTA/H₂O₂ is able to degrade metoprolol at initial neutral pH, and the degradation rate

of metoprolol can be greatly improved with the presence of Mn^{II}. The *k* values of metoprolol in Fe^{III}-NTA/H₂O₂-Mn system were typically 7–9-fold larger than those in Fe^{III}-NTA/H₂O₂ system. It can be concluded that the presence of Mn^{II} does not directly catalyze Fenton-like reaction, but indirectly enhance Fe^{III}-NTA-catalyzed Fenton-like reaction over a wide pH range of 4.0–8.0. The main intermediates during the degradation of metoprolol have been identified by mass spectrometry, which suggested that the degradation of metoprolol and its intermediates were mainly caused by HO[•] attack. Fe^{III}-NTA/H₂O₂-Mn system led to more efficient degradation of metoprolol and its intermediates.

Acknowledgments

This work was supported in part by the Innovation Scientists and Technicians Troop Construction Projects of Henan Province, Plan for Scientific Innovation Talent of Henan Province (Grant No. 134200510014), and in part by the Natural Science Foundation of Jiangsu Province (Grant No. BK20140341). The authors also would like to thank anonymous reviewers for their thoughtful comments and suggestions on this manuscript.

Appendix A. Supplementary data

Quenching tests with 2-propanol and chloroform on the degradation of metoprolol in Fe^{III}-NTA/H₂O₂-Mn system (Fig. S1). Total molecular ion chromatogram of metoprolol and its intermediates in Fe^{III}-NTA/H₂O₂ and Fe^{III}-NTA/H₂O₂-Mn systems (Fig. S2 and Fig. S3). Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.cej.2016.12.098>.

References

- [1] M.S. Kostich, A.L. Batt, J.M. Lazorchak, Concentrations of prioritized pharmaceuticals in effluents from 50 large wastewater treatment plants in the US and implications for risk estimation, *Environ. Pollut.* 184 (2014) 354–359.
- [2] J. Sun, Q. Luo, D. Wang, Z. Wang, Occurrences of pharmaceuticals in drinking water sources of major river watersheds, China, *Ecotoxicol. Environ. Saf.* 117 (2015) 132–140.
- [3] Q.-T. Liu, H.E. Williams, Kinetics and degradation products for direct photolysis of β-blockers in water, *Environ. Sci. Technol.* 41 (2007) 803–810.
- [4] C. Miège, J.M. Choubert, L. Ribeiro, M. Eusèbe, M. Coquery, Fate of pharmaceuticals and personal care products in wastewater treatment plants – conception of a database and first results, *Environ. Pollut.* 157 (2009) 1721–1726.
- [5] C.-C. Chen, R.-J. Wu, Y.-Y. Tzeng, C.-S. Lu, Chemical oxidative degradation of acridine orange dye in aqueous solution by Fenton's reagent, *J. Chin. Chem. Soc.* 56 (2009) 1147–1155.
- [6] H.-J. Fan, S.-T. Huang, W.-H. Chung, J.-L. Jan, W.-Y. Lin, C.-C. Chen, Degradation pathways of crystal violet by Fenton and Fenton-like systems: condition optimization and intermediate separation and identification, *J. Hazard. Mater.* 171 (2009) 1032–1044.
- [7] C.-C. Chen, W.-C. Chen, M.-R. Chiou, S.-W. Chen, Y.Y. Chen, H.-J. Fan, Degradation of crystal violet by an FeGAC/H₂O₂ process, *J. Hazard. Mater.* 196 (2011) 420–425.
- [8] B. Abramović, S. Kler, D. Šojić, M. Laušević, T. Radović, D. Vione, Photocatalytic degradation of metoprolol tartrate in suspensions of two TiO₂-based photocatalysts with different surface area. Identification of intermediates and proposal of degradation pathways, *J. Hazard. Mater.* 198 (2011) 123–132.
- [9] F.J. Rivas, O. Gimeno, T. Borralho, M. Carbajo, UV-C radiation based methods for aqueous metoprolol elimination, *J. Hazard. Mater.* 179 (2010) 357–362.
- [10] V. Romero, S. Acevedo, P. Marco, J. Giménez, S. Esplugas, Enhancement of Fenton and photo-Fenton processes at initial circumneutral pH for the degradation of the β-blocker metoprolol, *Water Res.* 88 (2016) 449–457.
- [11] Y. Sun, J.J. Pignatello, Chemical treatment of pesticide wastes. Evaluation of iron(III) chelates for catalytic hydrogen peroxide oxidation of 2,4-D at circumneutral pH, *J. Agric. Food Chem.* 40 (1992) 322–327.
- [12] A. Rastogi, S.R. Al-Abed, D.D. Dionysiou, Effect of inorganic, synthetic and naturally occurring chelating agents on Fe(II) mediated advanced oxidation of chlorophenols, *Water Res.* 43 (2009) 684–694.
- [13] W. Huang, M. Brigante, F. Wu, C. Mousty, K. Hanna, G. Mailhot, Assessment of the Fe(III)-EDDS complex in Fenton-like processes: from the radical formation to the degradation of bisphenol A, *Environ. Sci. Technol.* 47 (2013) 1952–1959.

- [14] Y. Wu, M. Passananti, M. Brigante, W. Dong, G. Mailhot, Fe(III)–EDDS complex in Fenton and photo-Fenton processes: from the radical formation to the degradation of a target compound, *Environ. Sci. Pollut. Res.* 21 (2014) 12154–12162.
- [15] Y. Zhang, N. Klammerth, S.A. Messele, P. Chelme-Ayala, M. Gamal El-Din, Kinetics study on the degradation of a model naphthenic acid by ethylenediamine-N, N'-disuccinic acid-modified Fenton process, *J. Hazard. Mater.* 318 (2016) 371–378.
- [16] N. Klammerth, S. Malato, A. Agüera, A. Fernández-Alba, G. Mailhot, Treatment of municipal wastewater treatment plant effluents with modified photo-Fenton as a tertiary treatment for the degradation of micro pollutants and disinfection, *Environ. Sci. Technol.* 46 (2012) 2885–2892.
- [17] N. Klammerth, S. Malato, A. Agüera, A. Fernández-Alba, Photo-Fenton and modified photo-Fenton at neutral pH for the treatment of emerging contaminants in wastewater treatment plant effluents: a comparison, *Water Res.* 47 (2013) 833–840.
- [18] S. Miralles-Cuevas, I. Oller, J.A.S. Pérez, S. Malato, Removal of pharmaceuticals from MWTP effluent by nanofiltration and solar photo-Fenton using two different iron complexes at neutral pH, *Water Res.* 64 (2014) 23–31.
- [19] Y.H. Dao, J. De Laat, Hydroxyl radical involvement in the decomposition of hydrogen peroxide by ferrous and ferric-nitritotriacetate complexes at neutral pH, *Water Res.* 45 (2011) 3309–3317.
- [20] S.-P. Sun, X. Zeng, A.T. Lemley, Kinetics and mechanism of carbamazepine degradation by a modified Fenton-like reaction with ferric-nitritotriacetate complexes, *J. Hazard. Mater.* 252–253 (2013) 155–165.
- [21] Y. Zhang, N. Klammerth, M. Gamal El-Din, Degradation of a model naphthenic acid by nitritotriacetic acid – modified Fenton process, *Chem. Eng. J.* 292 (2016) 340–347.
- [22] A. De Luca, R.F. Dantas, S. Esplugas, Assessment of iron chelates efficiency for photo-Fenton at neutral pH, *Water Res.* 61 (2014) 232–242.
- [23] Y. Li, J. Sun, S.-P. Sun, Mn²⁺-mediated homogeneous Fenton-like reaction of Fe(III)-NTA complex for efficient degradation of organic contaminants under neutral conditions, *J. Hazard. Mater.* 313 (2016) 193–200.
- [24] C. Kormann, D.W. Bahnemann, M.R. Hoffmann, Photocatalytic production of hydrogen peroxides and organic peroxides in aqueous suspensions of titanium dioxide, zinc oxide, and desert sand, *Environ. Sci. Technol.* 22 (1988) 798–806.
- [25] R.J. Watts, J. Sarasa, F.J. Loge, A.L. Teel, Oxidative and reductive pathways in manganese-catalyzed Fenton's reactions, *J. Environ. Eng. ASCE* 131 (2005) 158–164.
- [26] B.H.J. Bielski, D.E. Cabelli, R.L. Arudi, A.B. Ross, Reactivity of HO₂/O₂⁻ radicals in aqueous solution, *J. Phys. Chem. Ref. Data* 14 (1985) 1041–1100.
- [27] J. De Laat, Y.H. Dao, N. Hamdi El Najjar, C. Daou, Effect of some parameters on the rate of the catalysed decomposition of hydrogen peroxide by iron(III)-nitritotriacetate in water, *Water Res.* 45 (2011) 5654–5664.
- [28] G.V. Buxton, C.L. Greenstock, W.P. Helman, A.B. Ross, Critical-review of rate constants for reactions of hydrated electrons, hydrogen-atoms and hydroxyl radicals (·OH/O⁻) in aqueous-solution, *J. Phys. Chem. Ref. Data* 17 (1988) 513–886.
- [29] R.J. Motekaitis, A.E. Martell, The iron(III) and iron(II) complexes of nitritotriacetic acid, *J. Coord. Chem.* 31 (1994) 67–78.
- [30] B.A. Smith, A.L. Teel, R.J. Watts, Identification of the reactive oxygen species responsible for carbon tetrachloride degradation in modified Fenton's systems, *Environ. Sci. Technol.* 38 (2004) 5465–5469.
- [31] M. Bucheli-Witschel, T. Egli, Environmental fate and microbial degradation of aminopolycarboxylic acids, *FEMS Microbiol. Rev.* 25 (2001) 69–106.
- [32] C. Slegers, G. Baldacchino, D. Le Parc, B. Hickel, B. Tilquin, Radical mechanisms in the radiosterilization of metoprolol tartrate solutions, *Pharm. Res.* 20 (2003) 1977–1983.
- [33] W. Song, W.J. Cooper, S.P. Mezyk, J. Greaves, B.M. Peake, Free radical destruction of β-blockers in aqueous solution, *Environ. Sci. Technol.* 42 (2008) 1256–1261.
- [34] C. Walling, Fenton's reagent revisited, *Acc. Chem. Res.* 8 (1975) 125–131.
- [35] H. Yang, T. An, G. Li, W. Song, W.J. Cooper, H. Luo, X. Guo, Photocatalytic degradation kinetics and mechanism of environmental pharmaceuticals in aqueous suspension of TiO₂: a case of β-blockers, *J. Hazard. Mater.* 179 (2010) 834–839.
- [36] J. Radjenovic, B.I. Escher, K. Rabaey, Electrochemical degradation of the β-blocker metoprolol by Ti/Ru_{0.7}Ir_{0.3}O₂ and Ti/SnO₂-Sb electrodes, *Water Res.* 45 (2011) 3205–3214.